


LITERATURA

1. Chapple C, Kelleher C, Siddiqui E, et al. Validation of the Overactive Bladder-Bladder Assessment Tool (OAB-BAT): A Potential Alternative to the Standard Bladder Diary for Monitoring OAB Outcomes. *Eur Urol Focus*. 2021;7(5):1176-1183.
2. Malde S, Kelly S, Saad S, et al. Case-finding tools for the diagnosis of OAB in women: A narrative review. *Neurourol Urodyn*. 2020;39(1):13-24.
3. Rosier PFWM, Schaefer W, Lose G, et al. International Continence Society Good Urodynamic Practices and Terms 2016: Urodynamics, uroflowmetry, cystometry, and pressure-flow study. *Neurourol Urodyn*. 2017;36(5):1243-1260.
4. Brown JS, McNaughton KS, Wyman JF, et al. Measurement characteristics of a voiding diary for use by men and women with overactive bladder. *Urology*. 2003;61(4):802-809.
5. Lukacz ES, DuHamel E, Menefee SA, et al. Elevated postvoid residual in women with pelvic floor disorders: prevalence and associated risk factors. *Int Urogynecol J Pelvic Floor Dysfunct*. 2007;18(4):397-400.
6. Siroky MB, Olsson CA, Krane RJ. The flow rate nomogram: I. Development. *J Urol*. 1979;122(5):665-668.
7. Working Group of the United Kingdom Continence Society; Abrams P, Eustice S, Gammie A, et al. United Kingdom Continence Society: Minimum standards for urodynamic studies, 2018. *Neurourol Urodyn*. 2019;38(2):838-856.
8. Dörflinger A, Gorton E, Stanton S, Dreher E. Urethral pressure profile: is it affected by position? *Neurourol Urodyn*. 2002;21(6):553-557.
9. Emberton M. Definition of at-risk patients: dynamic variables. *BJU Int*. 2006;97 Suppl 2:12-15.
10. Mochtar CA, Kiemeny LA, van Riemsdijk MM, et al. Post-void residual urine volume is not a good predictor of the need for invasive therapy among patients with benign prostatic hyperplasia. *J Urol*. 2006;175(1):213-216.
11. Abrams P. Bladder outlet obstruction index, bladder contractility index and bladder voiding efficiency: three simple indices to define bladder voiding function. *BJU Int*. 1999;84(1):14-15.
12. Nitti VW. Pressure flow urodynamic studies: the gold standard for diagnosing bladder outlet obstruction. *Rev Urol*. 2005;7(Suppl 6):S14-21.
13. Siroky MB, Olsson CA, Krane RJ. The flow rate nomogram: II. Clinical correlation. *J Urol*. 1980;123(2):208-210.
14. Reynard JM, Peters TJ, Lim C, et al. The value of multiple free-flow studies in men with lower urinary tract symptoms. *Br J Urol*. 1996;77(6):813-818.
15. Abrams P, Cardozo L, Fall M, et al. The standardisation of terminology of lower urinary tract function: report from the Standardisation Sub-committee of the International Continence Society. *Neurourol Urodyn*. 2002;21(2):167-178.
16. Jeong SJ, Kim HJ, Lee YJ, et al. Prevalence and Clinical Features of Detrusor Underactivity among Elderly with Lower Urinary Tract Symptoms: A Comparison between Men and Women. *Korean J Urol*. 2012;53(5):342-348.
17. Thomas AW, Cannon A, Bartlett E, et al. The natural history of lower urinary tract dysfunction in men: the influence of detrusor underactivity on the outcome after transurethral resection of the prostate with a minimum 10-year urodynamic follow-up. *BJU Int*. 2004;93(6):745-750.
18. Al-Hayek S, Thomas A, Abrams P. Natural history of detrusor contractility--minimum ten-year urodynamic follow-up in men with bladder outlet obstruction and those with detrusor. *Scand J Urol Nephrol Suppl*. 2004;(215):101-108.
19. Thomas AW, Cannon A, Bartlett E, et al. The natural history of lower urinary tract dysfunction in men: minimum 10-year urodynamic followup of transurethral resection of prostate for bladder outlet obstruction. *J Urol*. 2005;174(5):1887-1891.

Právě jste dokončili studium medicíny?

Máme pro vás odborný kongres zcela zdarma.



SOLEN MEDICAL EDUCATION

